

Triangular hyperbolic buildings.

Riikka Kangaslampi, Alina Vdovina

February 1, 2008

Abstract

We construct triangular hyperbolic polyhedra whose links are generalized 4-gons. The universal cover of those polyhedra are hyperbolic buildings, which apartments are hyperbolic planes tessellated by regular triangles with angles $\pi/4$. Moreover, the fundamental groups of the polyhedra acts simply transitively on vertices of the buildings.

Introduction

A *polyhedron* is a two-dimensional complex which is obtained from several oriented p -gons by identification of corresponding sides. We will consider euclidean and hyperbolic polyhedra. Let's take a point of the polyhedron and take a sphere (euclidean or hyperbolic) of a small radius at this point. The intersection of the sphere with the polyhedron is a graph, which is called the *link* at this point.

We consider polyhedra such that all links of all vertices are generalized 4-gons.

We say that a polyhedron P is an (m, n) -polyhedron if the girth of any link of P is at least m and each face of P is a polygon with at least n edges.

Let P be a (m, n) -polyhedron such that m and n satisfy the inequality $mn \geq 2(m + n)$. This inequality appears, for example, in small cancellation theory [13]. The universal covering of a (m, n) -polyhedron with the metric introduced in [1, p. 165] is a complete metric space of non-positive curvature in the sense of Alexandrov and Busemann [11].

In this note we construct examples of $(4, 3)$ -polyhedra. It follows from [3], that the fundamental groups of our polyhedra satisfy the property (T) of Kazhdan. (Another relevant reference is [16])

Definition. A *generalized m -gon* is a graph which is a union of subgraphs, called apartments, such that:

1. Every apartment is a cycle composed of $2m$ edges for some fixed m .
2. For any two edges there is an apartment containing both of them.
3. If two apartments have a common edge, then there is an isomorphism between them fixing their intersection pointwise.

Definition. Let $\mathcal{P}(p, m)$ be a tessellation of the hyperbolic plane by regular polygons with p sides, with angles π/m in each vertex where m is an integer. A *hyperbolic building* is a polygonal complex X , which can be expressed as the union of subcomplexes called apartments such that:

1. Every apartment is isomorphic to $\mathcal{P}(p, m)$.
2. For any two polygons of X , there is an apartment containing both of them.
3. For any two apartments $A_1, A_2 \in X$ containing the same polygon, there exists an isomorphism $A_1 \rightarrow A_2$ fixing $A_1 \cap A_2$.

Let C_p be a polyhedron whose faces are p -gons and whose links are generalized m -gons with $mp > 2m + p$. We equip every face of C_p with the hyperbolic metric such that all sides of the polygons are geodesics and all angles are π/m . Then the universal covering of such polyhedron is a hyperbolic building, see [9]. The polyhedra we construct in this note are from this class, since $p = 3$, $m = 4$. Our construction gives examples of hyperbolic triangular buildings with regular triangles as chambers, which were known before, on our knowledge. Examples of hyperbolic buildings with right-angled triangles were constructed by M.Bourdon [7].

In the case $p = 3$, $m = 3$, i.e. C_p is a simplex, we can give a euclidean metric to every face. In this metric all sides of the triangles are geodesics. The universal coverings of these polyhedra with the euclidean metric are euclidean buildings, see [2], [4].

Recall (cf. [14]) that a *generalized m -gon* is a connected, bipartite graph of diameter m and girth $2m$, in which each vertex lies on at least two edges. A graph is *bipartite* if its set of vertices can be partitioned into two disjoint subsets such that no two vertices in the same subset lie on a common edge. The vertices of one subset we will call black vertices and the vertices of the other subset the white ones. The *diameter* is the maximum distance between two vertices and the *girth* is the length of a shortest circuit.

Let G be a connected bipartite graph on $q + r$ vertices, q black vertices and r white ones. Let $\mathcal{A}$ and $\mathcal{B}$ be two alphabets on q and r letters respectively, $\mathcal{A} = \{x_1, x_2, \dots, x_q\}$ and $\mathcal{B} = \{y_1, y_2, \dots, y_r\}$. We mark every black vertex with an element from $\mathcal{A}$ and every white vertex with an element from $\mathcal{B}$.

1 Polygonal presentation.

We recall a definition of polygonal presentation introduced in [15].

Definition. Suppose we have n disjoint connected bipartite graphs $G_1, G_2, \dots, G_n$. Let P_i and Q_i be the sets of black and white vertices respectively in G_i , $i = 1, \dots, n$; let $P = \bigcup P_i$, $Q = \bigcup Q_i$, $P_i \cap P_j = \emptyset$, $Q_i \cap Q_j = \emptyset$ for $i \neq j$ and let λ be a bijection $\lambda: P \rightarrow Q$.

A set $\mathcal{K}$ of k -tuples $(x_1, x_2, \dots, x_k)$, $x_i \in P$, will be called a *polygonal presentation* over P compatible with λ if

- (1) $(x_1, x_2, x_3, \dots, x_k) \in \mathcal{K}$ implies that $(x_2, x_3, \dots, x_k, x_1) \in \mathcal{K}$;
- (2) given $x_1, x_2 \in P$, then $(x_1, x_2, x_3, \dots, x_k) \in \mathcal{K}$ for some $x_3, \dots, x_k$ if and only if x_2 and $\lambda(x_1)$ are incident in some G_i ;
- (3) given $x_1, x_2 \in P$, then $(x_1, x_2, x_3, \dots, x_k) \in \mathcal{K}$ for at most one $x_3 \in P$.

If there exists such $\mathcal{K}$, we will call λ a *basic bijection*.

Polygonal presentations for $n = 1$, $k = 3$ were listed in [8] with the incidence graph of the finite projective plane of order two or three as the graph G_1 .

Some polygonal presentations for $k > 3$ were constructed in [15].

Now we construct polygonal presentations with $k = 3, n = 1$, but the graph G_1 is a generalized 4-gon.

T_1 :

(x_1, x_2, x_7)
 (x_1, x_8, x_{11})
 (x_1, x_{14}, x_5)
 (x_2, x_4, x_{13})
 (x_{12}, x_4, x_2)
 (x_4, x_9, x_3)
 (x_6, x_8, x_3)
 (x_{14}, x_6, x_3)
 (x_{12}, x_{10}, x_5)
 (x_{13}, x_{15}, x_5)
 (x_{12}, x_9, x_6)
 (x_{11}, x_{10}, x_7)
 (x_{14}, x_{13}, x_7)
 (x_9, x_{15}, x_8)
 (x_{11}, x_{15}, x_{10})

T_2 :

(x_1, x_{10}, x_1)
 (x_1, x_{15}, x_2)
 (x_2, x_{11}, x_9)
 (x_2, x_{14}, x_3)
 (x_3, x_7, x_4)
 (x_3, x_{15}, x_{13})
 (x_4, x_8, x_6)
 (x_{12}, x_{11}, x_4)
 (x_5, x_8, x_5)
 (x_5, x_{10}, x_{12})
 (x_6, x_{14}, x_6)
 (x_7, x_{12}, x_7)
 (x_{13}, x_9, x_8)
 (x_{14}, x_{15}, x_9)
 (x_{13}, x_{11}, x_{10})

Let's show, that these sets are desired polygonal presentations. Remark, that the smallest generalized 4-gon can be presented in the following way: its "points" are pairs (ij) , where $i, j = 1, \dots, 6$, $i \neq j$ and "lines" are

triples of those pairs where all i, j are different. We mark pairs (ij) , where $i, j = 1, \dots, 6$, $i \neq j$ by numbers from 1 to 15 in natural order. Now one can check by direct examination, that the graph G_1 is really the smallest generalized 4-gon. The polygonal presentation T_1 and T_2 are not equivalent since there is no automorphism of the generalized 4-gon, which transforms one to another.

2 Construction of polyhedra.

We can associate a polyhedron K on n vertices with each polygonal presentation $\mathcal{K}$ as follows: for every cyclic k -tuple $(x_1, x_2, x_3, \dots, x_k)$ from the definition we take an oriented k -gon on the boundary of which the word $x_1 x_2 x_3 \dots x_k$ is written. To obtain the polyhedron we identify the corresponding sides of our polygons, respecting orientation. We will say that the polyhedron K *corresponds* to the polygonal presentation $\mathcal{K}$.

Lemma [15] A polyhedron K which corresponds to a polygonal presentation $\mathcal{K}$ has graphs $G_1, G_2, \dots, G_n$ as the links.

Remark. Consider a polygonal presentation $\mathcal{K}$. Let s_i be the number of vertices of the graph G_i and t_i be the number of edges of G_i , $i = 1, \dots, n$. If the polyhedron K corresponds to the polygonal presentation $\mathcal{K}$, then K has n vertices (the number of vertices of K is equal to the number of graphs), $k \sum_{i=1}^n s_i$ edges and $\sum_{i=1}^n t_i$ faces, all faces are polygons with k sides.

For polygonal presentation T_i , $i = 1, 2$ take 15 oriented regular hyperbolic triangles with angles $\pi/4$, write words from the presentation on their boundaries and glue together respecting orientation sides with the same letters. The result is a hyperbolic polyhedron with one vertex and 15 faces and the universal covering is a triangular hyperbolic building. The fundamental group of the polyhedron acts simply transitively on vertices of the building. The group has 15 generators and 15 relations, which come naturally from the polygonal presentation.

References

- [1] W. Ballmann, M. Brin, *Polygonal complexes and combinatorial group theory*, Geometriae Dedicata 50 (1994), 165–191.
- [2] W. Ballmann, M. Brin, *Orbihedra of nonpositive curvature*, Publications Mathématiques IHES, 82 (1995), 169–209.
- [3] W. Ballmann, J. Świątkowski, *On L^2 -cohomology and property (T) for automorphism groups of polyhedral cell complexes*, Geom.Funct.Anal. 7 (1997), no.4, 615–645.
- [4] S. Barre, *Polyèdres finis de dimension 2 à courbure ≤ 0 et de rang 2*, Ann. Inst. Fourier 45, 4 (1995), 1037–1059.
- [5] M. Bourdon, *Immeubles hyperboliques, dimension conforme et rigidité de Mostow*, Geom. Funct. Anal. 7 (1997), 245–268.

- [6] M. Bourdon, *Sur la dimension de Hausdorff de l'ensemble limite de sous-groupes convexes co-compacts*, C. R. Acad. Sci. Paris, Série 1, 325 (1997), 1097–1100.
- [7] M. Bourdon, *Sur les immeubles fuchsien et leur type de quasi-isométrie*, Ergodic Theory Dynam. Systems, 20 (2000), no. 2, 343–364.
- [8] D. Cartwright, A. Mantero, T. Steger, A. Zappa, *Groups acting simply transitively on vertices of a building of type $\tilde{A}_2$* , Geometriae Dedicata 47 (1993), 143–166.
- [9] D. Gaboriau, F. Paulin, *Sur les immeubles hyperboliques*, Geometriae Dedicata 88 (2001), no. 1–3, 153–197.
- [10] R. Charney, A. Lytchak, *Metric characterization of spherical and Euclidean buildings*, Geom. Topol. 5 (2001), 521–550 (electronic).
- [11] E. Ghys, P. de la Harpe (eds.), *Sur les groupes Hyperboliques d'après Mikhael Gromov*, Birkhäuser, Boston, Basel, Berlin, 1990.
- [12] M. Gromov, *Hyperbolic groups*, Essays in Group Theory (ed. M. Gersten) MSRI Publ. 8, Springer, 1987, 75–263.
- [13] R. Lyndon, P. Schupp, *Combinatorial group theory*, Springer, Berlin, Heidelberg, New York, 1977.
- [14] M. Ronan, *Lectures on buildings*, Perspectives in mathematics; vol. 7 (1989).
- [15] A. Vdovina, *Combinatorial structure of some hyperbolic buildings*, Math. Z. 241 (2002), no. 3, 471–478.
- [16] A. Zuk, *La propriété de Kazhdan pour les groupes agissant sur les polyèdres*, C.R.A.S. Sér.1 Math(1996), no.5, 453–458.